arXiv:hep-ph/0607212v1 19 Jul 2006

Comment on "Mass and KΛ Coupling of the N*(1535)"

A. Sibirtsev[1], J. Haidenbauer[2], and Ulf-G. Meißner,[1,2]
[1] *HISK, Universität Bonn, Nußallee 14-16, D-53115 Bonn, Germany and*
[2] *Institut für Kernphysik (Theorie), Forschungszentrum Jülich, D-52425 Jülich, Germany*

Recently it has been argued that the effective coupling of the $N^*(1535)$ to $K\Lambda$ is even larger than its coupling to the ηN channel [1] from an analysis of data on the $J/\Psi \to \bar{p}\Lambda K$ and $J/\Psi \to \bar{p}p\eta$ decays within a resonance isobar model. Based on those couplings of $N^*(1535)$ to the ηN and $K\Lambda$ (and the πN) states new properties of the resonance have been derived, namely a mass of 1400 MeV and a width of 270 MeV, that differ radically from the standard values.

As additional if not decisive argument in favor of the strong $N^*(1535)$ coupling to the $K\Lambda$ channel the authors of Ref. [1] claimed that the low-energy cross section data on the reaction $pp \to pK^+\Lambda$ are compatible with a large contribution from the $N^*(1535)$ resonance and that it is, in fact, needed to reproduce the experimental result. In their calculation based on tree-level Feynman diagrams the final-state interaction (FSI) between the proton and the Λ was neglected and it was concluded that the $N^*(1535)$ dominates the reaction at low energies.

There are several points in the procedure followed in Ref. [1] that one might criticize. For example, why was only the $N^*(1535)$ resonance included in the analysis of the $\bar{p}K\Lambda$ decay branching ratio, but not the $N^*(1650)$? The latter choice seems much more natural since this resonance is much closer to the threshold of the $J/\Psi \to \bar{p}K\Lambda$ decay. However, in our comment we want to concentrate on their results for $pp \to pK^+\Lambda$. First, let us mention that their calculation is in contradiction with all other interpretations of the $pp \to pK^+\Lambda$ cross section where the enhancement at low energies is due to the Λp FSI rather than the $N^*(1535)$ resonance [2, 3, 4]. The result in Ref. [1] is also in disagreement with the experimental Dalitz plot [5] which strongly suggests that at low energies the $pp \to pK^+\Lambda$ reaction is dominated by the contribution from the $N^*(1650)$ resonance.

In Ref. [4] we proposed to study the role of the FSI in the $pp \to pK^+\Lambda$ reaction by analyzing differential observables, namely the Dalitz plot distribution. By making cuts on the $K^+\Lambda$ invariant mass one can isolate the contribution from N^* resonances and project the Dalitz plot on the Λp axis. In that case an enhancement at low Λp masses would manifest the presence of a FSI. It is clear [4] that N^* resonances that couple to the $K^+\Lambda$ system can never produce such an enhancement.

Recently the COSY-TOF Collaboration [5] measured the Dalitz plot distribution and presented the projections on the Λp invariant mass with different cuts on the $K^+\Lambda$ masses. The Λp spectra are shown in Ref. [5] in terms of the angular distributions of the Λ in reference to the proton direction in the $K^+\Lambda$ cm system, i.e. as function of the Λp helicity angle θ_H. The relation between θ_H and the Λp invariant mass is given by Eq. (8) of our paper [4]. Fig. 1 shows experimental results for the cut on the $K^+\Lambda$ mass chosen in order to isolate the contribution from low lying N^* resonances (with the convention $-\theta_H$ applied in the experiment [5] so that the lowest Λp masses corresponds to $\cos\theta_H$=1). The data indicate a strong θ_H asymmetry. The solid histogram is our result obtained with inclusion of the Λp FSI. The dashed line shows a calculation employing the $N^*(1535)$ parameters claimed in Ref. [1] but without FSI. It is obvious that the latter scenario is in disagreement with the experiment.



FIG. 1: The Λp helicity angle spectra for $K^+\Lambda$ masses $M_{K\Lambda}$>1.74 GeV for the reaction $pp \to pK^+\Lambda$ at p_{beam} = 2.85 GeV/c. The solid histogram is our calculation including the Λp FSI, while the dashed line is obtained without FSI but with the $N^*(1535)$ parameters from Ref. [1]. The data are from Ref. [5].

We consider the experimental observation of Ref. [5] as a direct evidence for the presence of the Λp FSI. Thus, the results for $pp \to pK^+\Lambda$ by Liu and Zou [1] are rather questionable, and consequently also their conclusions based on the large $N^*(1535)$ coupling to the $K\Lambda$ system.

[1] B.C. Liu, and B.S. Zou, Phys. Rev. Lett. **96**, 042002 (2006).
[2] P. Moskal et al., Prog. Part. Nucl. Phys. **49**, 1 (2000).
[3] R. Shyam et al., Phys. Rev. C **63**, 022202 (2001).
[4] A. Sibirtsev et al., Eur. Phys. J. A **27**, 269 (2006).
[5] S. Abd El-Samad *et al.*, Phys. Lett. B **632**, 27 (2006).